

Five Things Physicians and Patients Should Question

1 Don't order a routine bone scan and CT scan of the pelvis in men with low-risk prostate cancer.

Low-risk patients (defined by D'Amico criteria and National Comprehensive Cancer Network guidelines) are unlikely to have metastatic disease. Accordingly, imaging is generally unnecessary in patients with newly diagnosed prostate cancer who have a PSA <20.0 ng/mL and a Gleason score 6 or less unless the patient's history or clinical examination suggests distant disease. Metastases are much more likely in high-grade disease that is characterized by fast and aggressive growth into surrounding areas such as bones or lymph nodes.

2 Don't order serum testosterone in men without symptoms of hypogonadism.

In the evaluation of men with erectile dysfunction, testosterone should only be ordered if there are signs and/or symptoms of hypogonadism.

3 Don't prescribe testosterone to men with erectile dysfunction who have normal testosterone levels.

While testosterone treatment may increase sexual interest, there appears to be no significant influence on erectile function in men with normal testosterone levels.

4 Don't use antimicrobials to treat asymptomatic bacteriuria in the elderly.

Studies suggest that asymptomatic bacteriuria in the elderly does not carry significant risk of morbidity if left untreated. Antimicrobial treatment studies for asymptomatic bacteriuria in older adults demonstrate no benefits and show increased adverse antimicrobial effects. Consensus criteria has been developed to characterize the specific clinical symptoms that, when associated with bacteriuria, define urinary tract infection. Screening for and treatment of asymptomatic bacteriuria is recommended before urologic procedures for which mucosal bleeding is anticipated.

5 Don't perform ultrasound on boys with cryptorchidism.

Ultrasound is of minimal value in localizing the position or existence of testes that cannot be felt through physical examination. Studies have shown that there remained a significant chance that testes were present even after a negative ultrasound result. The likelihood of locating testes is low when using ultrasound.

How the list was created

The Canadian Urological Association (CUA) utilized its guidelines committee to review evidence from the association's guidelines and the suggestions put forward by the American Urological Association (AUA) to identify potential topics for nomination to the CUA's *Choosing Wisely Canada* list. The general membership had the opportunity to put forward suggestions after being notified of the process via email. The committee reviewed a number of recommendations and through a consensus process identified the five tests or procedures that should be questioned. These recommendations were reviewed and approved by the CUA guidelines committee. Recommendations 1, 3 and 5 were adopted with permission from the Five Things Physicians and Patients Should Question. © 2013 American Urological Association. Recommendation 4 was adopted with permission from the American Geriatrics Society's, "Ten Things Physicians and Patients Should Question"©; The American Geriatrics Society, New York, New York. Reprinted with permission. www.americangeriatrics.org.

Sources

- 1** American Urological Association. Prostate-Specific Antigen Best Practice Statement [Internet]. Linthicum (MD): AUA; 2009 [cited 2012 Oct 16]. Available from: www.auanet.org/content/guidelines-and-quality-care/clinical-guidelines/main-reports/psa09.pdf.
National Comprehensive Cancer Network. National Comprehensive Cancer Network clinical practice guidelines in oncology (NCCN Guidelines®): prostate cancer. Fort Washington (PA): NCCN; 2012 Apr.
- 2** American Urological Association. Management of Erectile Dysfunction Clinical Practice Guideline [Internet]. Linthicum (MD): AUA; 2005 [updated 2007; cited 2012 Oct 16]. Available from: <http://www.auanet.org/common/pdf/education/clinical-guidance/Erectile-Dysfunction.pdf>
The Endocrine Society. Testosterone Therapy in Adult Men with Androgen Deficiency Syndromes: An Endocrinology Society Clinical Practice Guideline [Internet]. 2010 [cited 2014 Sep 22]. Available from: <http://www.endocrine.org/~media/endosociety/Files/Publications/Clinical%20Practice%20Guidelines/FINAL-Androgens-in-Men-Standalone.pdf>
- 3** American Urological Association. Management of Erectile Dysfunction Clinical Practice Guideline [Internet]. Linthicum (MD): AUA; 2005 [updated 2007; cited 2012 Oct 16]. Available from: <http://www.auanet.org/common/pdf/education/clinical-guidance/Erectile-Dysfunction.pdf>
- 4** Juthani-Mehta M. Asymptomatic bacteriuria and urinary tract infection in older adults. *Clin Geriatr Med*. 2007 Aug;23(3):585–594.
Nordenstam GR, Brandberg CA, Odén AS, Svanborg Edén CM, Svanborg A. Bacteriuria and mortality in an elderly population. *N Engl J Med*. 1986 May 1;314(18):1152–1156.
Nicolle LE, Mayhew WJ, Bryan L. Prospective randomized comparison of therapy and no therapy for asymptomatic bacteriuria in institutionalized elderly women. *Am J Med*. 1987 Jul;83(1):27–33.
Nicolle LE, Bradley S, Colgan R, Rice JC, Schaeffer A, Hooton TM; Infectious Diseases Society of America; American Society of Nephrology; American Geriatric Society. Infectious Diseases Society of America Guidelines for the diagnosis and treatment of asymptomatic bacteriuria in adults. *Clin Infect Dis*. 2005 Mar 1;40(5):643–65.
- 5** Tasian G and Copp H: Diagnostic performance of ultrasound in Nonpalpable Cryptorchidism: A systematic review and meta-analysis. *Pediatrics* . 2011 Jan: 127(1): 119–128.
National Advisory Committee on Blood and Blood Products (NAC). [Internet]. [cited 2014 May 16]. NAC PCC Working Group. Recommendations for Use of Prothrombin Complex Concentrates in Canada. Available from: <http://www.nacblood.ca/resources/guidelines/PCC.html>
Scottish Intercollegiate Guidelines Network (SIGN). Antithrombotics: indications and management. Edinburgh (UK): 2012. 75 p. Report No. 129.

About Choosing Wisely Canada

Choosing Wisely Canada is a campaign to help physicians and patients engage in conversations about unnecessary tests, treatments and procedures, and to help physicians and patients make smart and effective choices to ensure high-quality care.

For more information on *Choosing Wisely Canada* or to see other lists of Five Things Physicians and Patients Should Question, visit www.choosingwiselycanada.org. Join the conversation on Twitter @ChooseWiselyCA.

About The Canadian Urological Association

The Canadian Urological Association (CUA) is a proud partner of the *Choosing Wisely Canada* campaign. CUA represents and provides a voice for all Canadian urologists and fosters dedication of all members of the profession toward ensuring the highest possible standard of urologic care of Canadians.